



Northern Inland Region of California

This product is an output of the Climate-Informed Reforestation Workshop, hosted by the USDA California Climate Hub and funded by the California Department of Forestry and Fire Protection.

Regional experts contributed to the workshop and the content found in this guidance.

Editors

Aviv Karasov-Olson, Jessica DellaRossa, Jennifer Smith, Steven Ostoja

Contributors

Kyle Merriam, Ricky Satomi, Gabrielle Bohlman, Eric Knapp, Sophia Lemmo, Brian Lindstrand, Keli McElroy, Gary Urdahl

Graphic Design

Dudek Consulting Visual Storytelling Team Amelia Oxarart







Climate-Informed Reforestation Guidance For Red Fir Forests in Northern Inland California

Introduction to Climate-Informed Reforestation

Reforesting forests so they are resilient to future disturbances requires strategic planning, deliberate implementation, and follow-up management. In the face of climate change, those requirements are even more imperative. Changes in temperature, precipitation, and drought may push many species out of their bioclimatic envelope into environments that are more stressful, which can reduce health, survival, and productivity of trees. Generally, this stress will be exacerbated at lower elevations and for species which prefer colder and wetter climates.

Developing a strategic plan to assess the broader landscape context within which reforestation is considered is crucial for prioritizing reforestation success (Meyer et al., 2021; North et al., 2019). Climate-informed reforestation depends on effective site preparation, selection of appropriate planting material, and planting in densities, locations, and arrangements that will produce the best short- and long-term outcomes. Reforestation to produce resilient forests not only requires effective action during the first year of planting, but also follow-up actions in the years and decades to follow (Fig. 1). Post-planting management includes repetitive treatments to maintain control of competing vegetation and encourage or maintain the desired species composition, densities, and arrangements identified during the planning process. In many instances a post-planting management goal may also be to introduce (or re-introduce) fire as an on-going component of the ecological system.

This guidance describes how site preparation, planting, and post-planting management actions can promote climate- and fire- resilient forests. This guidance was co-produced with regional foresters and ecologists through workshops and group meetings. It incorporates both peer-reviewed literature and place-based knowledge and aims to provide recommendations for all land ownership types (public and private) where possible.

Reforestation Process

1 Site Preparation	pg. 3
2 Selecting Species & Composition	pg. 7
3 Planting	pg. 9
4 Post-Planting Management	pg. 17

PUTTING IT TOGETHER: A REFORESTATION TIMELINE

Figure 1. Climate-informed reforestation is a long-term process. Key steps, including planting, vegetation control, reintroduction of fire, and monitoring, span up to 30 years. Specific activities and their timelines will reflect project and location-specific considerations.

YEAR O Seedlings Planted


After a site has been prepared, reforestation begins with the planting of tree seedlings across the burned landscape to kick-start recovery.

Seedlings are planted with variable spacing to mimic natural forest structure and enhance resilience.

Microsites are used as much as possible.

YEARS 2 - 5 Vegetation Control

Shrubs and other vegetation compete with young trees. Herbicide or mechanical treatment reduces surrounding vegetation to help seedlings grow.



Shrub & vegetation growth can outpace seedlings. Targeted treatment can help ensure that trees have enough sunlight and space to establish themselves.

YEARS 12 - 18

Beneficial Fire

Prescribed fire is first introduced to reduce surface fuels and support tree growth. Seedlings are now young trees, better able to withstand low-intensity burns.



Beneficial fires improve ecosystem health, recycle nutrients, reduce shrub cover & fuel loads, and increase the fire tolerance of young trees. Beneficial fire should be applied periodically.

YEARS 18 - 25+

Monitoring

A mixed-age forest is forming. Trees are taller, the structure is more complex, and ongoing monitoring ensures long-term success.



Monitoring reforested sites is a priority to track tree growth, survival, and fuel conditions over time.



Climate Change Considerations

Effectively removing wood biomass and managing competing vegetation are critical components of site preparation to offset the risks associated with drought and severe wildfire.

- **Drought resilience:** Where climate projections include an increased water deficit, site preparation should include actions to reduce plants that will compete with planted trees for scarce water resources. Increasing temperatures and drought stress can be offset with a focus on creating and maintaining protected microsites in locations where trees will be planted.
- **Wildfire resilience:** Fuels management should be prioritized in regions with a high likelihood of high-severity fire, including high-severity reburns.

Existing tools coupled with novel approaches for site preparation can achieve many climate-related goals. The particular methods (e.g., mechanical, burning, and/or some combination) selected to achieve these outcomes should be based on the operational and environmental context of the project, following existing guidance e.g., Stewart et al. (2021).

Key Actions

<u>Fuels Management</u>

Effective and timely wood biomass removal reduces the risk of severe fire and allows for safe planting and post-planting treatments. It is important to remove standing dead material (snags) and surface fuels to levels that reduce future high-severity fire. This involves managing 1 and 10 hour fuels, which cannot be effectively treated by hand or machine. Ideally, these fuels should be less than 3 to 10 tons per acre, depending on stand productivity, forest type, and fuel loads of the surrounding landscape (Scott & Burgan, 2005). Remaining fuels that are patchy and discreet in coverage can contribute to structural heterogeneity which is common in frequent, low severity fire systems and may reduce the risk of high-severity reburn (Lydersen et al., 2019; Lydersen & North, 2012). While extensive removal of wood biomass is critical for

reducing fire risk, it is also desirable to retain some woody elements on the landscape for ecological value and wildlife habitat.

- Reduce snag density to restore natural fire regimes and reduce the risk of high-severity reburn.
- Retain at least 3-5 snags per acre, spaced variably across the stand to create stand heterogeneity. Retain a combination of individual larger snags (>30 inches diameter at breast height (DBH)) and clumps of snags in well-protected microsites (Fig. 2) that are likely to ensure greater longevity and biological relevance compared to scattered dispersed snags. Additionally, in areas not considered important for strategic fuel treatments, retain snags on marginal and unproductive planting sites or sites where access is difficult for planting (e.g., steep slopes >30% and far from roads, structures, and infrastructure).
- Consider incorporating future fuel breaks into site preparation practices to include thinning around planting units, to protect the planted areas, possibly in the form of shaded fuel breaks¹, which can enhance wildfire resilience and protect the planted areas.
- Where it is deemed appropriate to reduce fuels prior to planting, remaining dead tree removal should be completed in the first 18-24 months following the disturbance, possibly up to 5 years for trees and sites with slower rates of decay. Timely dead tree removal and other fuels reduction treatments reduces the risk of high-severity reburn and improves access for subsequent reforestation actions including additional site preparation, planting, and post-planting management. This timeline is particularly influenced by how rapidly snags become unmerchantable, which may impede the economic feasibility of removing the material from the site.

<u>Managing Competing Vegetation</u>

Managing competing vegetation can be an important climate-informed practice, as it can reduce the impact of projected increases in water stress and fire activity. Site conditions and the existing plant species inform whether vegetation on the site will compete with the planted seedlings or not, and determine the intensity at which vegetation on a site needs to be managed. When vegetation is expected to compete with planted seedlings for limited resources, controlling that vegetation provides those trees with a competitive advantage for accessing soil moisture, which is a primary determinant of the successful establishment of young trees following planting. Vegetation management can also help reduce fuel loading and continuity to lower the potential for future high-severity fire, and allow for fire to re-enter a stand sooner without negatively impacting planted trees.

¹ Shaded fuel breaks, between 0.25 and 0.5 mi in width and treated by mechanical thinning & prescribed burning to reduce ladder (trees < 20 inches DBH) and surface fuels that increase the risk of crown fires, can be located near roads and ridgetops (Low et al. 2023).

However, not all vegetation may impact planted seedlings equally. Retaining a small to moderate amount of shrub and herb cover may create opportunities to provide shaded and protected microsites for young trees, which can increase seedling survival (Marshall et al., 2023), and play a facilitative role in postfire and highly stressful climatic conditions (Fig. 2). Many native shrubs, grasses, and forbs are important for maintaining soil moisture, fixing nitrogen, providing heat sinks for fires at certain times of the year, and promoting carbon sequestration.

Some managers recommend eliminating all competing vegetation that might inhibit success of planted trees, others recommend controlling and maintaining <25%, while others may opt for controlling and maintaining up to 50% shrub cover depending on the site conditions, shrub species, tree species, and management objectives.

Examples of Conditions Affecting Level of Shrub Control

- In xeric sites or during periods of extreme drought when shrub growth may be reduced, shrubs may provide more beneficial shade coverage for seedlings and allow for maintenance of slightly higher shrub cover. However, on mesic sites, the benefit of shade coverage is only seen when planting happens in the first 2 years following fire (Sorenson et al., 2025).
- Extensive growth and high cover of *Ceanothus*, bear clover, and manzanita may require more shrub control than other less aggressive shrub species based on their potential to outcompete planted seedlings and impact establishment.
- **White fir seedlings** may be more tolerant of high shade cover by shrubs compared to pines.
- Some shrub species (e.g., blue elderberry, California hazelnut, several species of
 gooseberries and currants) provide high value as cultural resources and wildlife
 habitat and can provide shading for conifer seedlings where they exist. These species
 also rarely dominate a stand and do not create many competition problems for young
 seedlings.
- Moderate to high levels of native grass or forb cover (>25 to 50%) could benefit conifer seedlings by providing shading or increased soil nutrients in both xeric and mesic sites.
- Following die-off from disease or drought, thorough monitoring will determine the need for and level of shrub control. Target (spot) treatment of shrub and competing vegetation where seedlings are to be planted in areas of high shrub cover (>60%) to create patchy shrub cover, and perhaps allow for a higher level of shrub cover than desired for post-fire reforestation.

Soil Management

An increase in high-severity fire and drought poses problems for soil health of forests. There are several suggested practices for minimizing negative impacts to soil health and soil moisture retention, though there is less consensus on the usefulness of these practices in every case and their applicability will vary based on specific soil properties.

- Experiment with retaining shrub cover to promote soil health (see benefits described in *Managing Competing Vegetation*).
- Experiment with biochar before planting or utilize biochar plugs when available at the nursey,
 particularly on xeric sites. Biochar can enhance water holding capacity, increase soil organic matter,
 nutrient retention and soil pH, which can increase the survival of planted seedlings (Marsh et al., 2023).
 It is likely more beneficial in low productivity or acidic soils and may have limited efficacy or no benefit
 in more productive sites.
- Experiment with mycorrhizal additions, when bareroots are not inoculated at the nursery. Mycorrhizal inoculation can improve the soil's physical characteristics, improve plant nutrient supply and lead to increased growth (Neuenkamp et al., 2019).
- Minimize disturbance to the duff, litter, and mineral layers after salvage logging to avoid shrub resprouting.
- **Following high-severity fire,** hydrophobic soils are often created. Breaking up this layer to minimize erosion and increase water infiltration can be beneficial, particularly on mesic sites.
 - Consider using contour tilling to address these issues as needed.
 - Only consider using deep ripping on in lower elevation stands where there were post-fire hydrophobicity effects on soil as this treatment can result in undesirable soil impacts (e.g., increase in invasive plant cover).
 - However, if soils are healthy post-fire, these intensive treatments can have negative impacts on soil health, such as degrading wildlife habitat and facilitating post-fire cheatgrass invasion.

Management of Surrounding Remnant Stand Following Disease-Driven Mortality

Disease driven mortality events necessitating reforestation are likely to increase in the face of climate change. These mortality events will require additional pre-planting preparation.

- Consider creating a species-specific buffer surrounding the planted area to ensure that the disease does
 not spread to newly planted trees. This buffer can be achieved by removing live host species that may
 have been exposed to the disease but are not yet showing signs of decline, and/or planting a buffer of
 non-host species between the remnant stand and core planted area. The width of the buffer is
 dependent upon the specific disease being addressed.
- Apply the necessary chemical or biological treatment to the area, when available.

Climate Change Considerations

Changes in temperature, precipitation, and drought may create environments that are more stressful and outside the bioclimatic envelope of many species, which can reduce health, survival, and growth. In this region, managers have noted such impacts in low-elevation pine stands as well as in higher elevation red fir stands. Decisions in a reforestation project that can buffer these risks include:

- Increasing the proportion of drought resistant species, particularly in xeric sites.
- Potentially shifting the relative proportion of each species, based on specific site characteristics, pre-disturbance stand health, and post-disturbance monitoring, as well as a desire to increase diversity of seedlings to reestablish on a site.
- Using a combination of local and climate-adapted seeds to increase genetic diversity and better provide for future climatic shifts.

Key Decisions

Seed Sourcing

Seed and seedling source selection may be limited by seed crop availability, regional availability, and access. However, when the option is available, there are several recommendations for seed sourcing. .

- Use planting as an opportunity to increase the genetic diversity within species to better provide for future climatic shifts.
- Use a mixture seed from the same seed zone and seed from lower elevations (500-1000 ft, 1-2 bands lower) or from further south in the range should be used. Seeds from lower and more southernly elevation bands have been shown to be more tolerant of hotter and drier conditions projected to be more extreme in coming decades. Currently a 50% mix of local and a 50% mix of seedlings for lower elevations or further south are recommended. In these cases, post-planting monitoring should include tracking outcomes of different seed sources.
- Use a seed lot selection tool (e.g., <u>CAST</u> tool) and/or work with genetics experts for selecting non-local seed.

• Use seed from seed orchards where available for species as they generally have a broader band of elevations and climactic conditions that they are suitable for due to inherent genetic diversity in the orchards.

Species Selection & Composition

In the red fir forests of this region, experts recommend replanting red fir after a disturbance. If red fir are unlikely to persist - in areas of high climate exposure or where stands are threatened or declining due to a variety of stressors, including drought, insects, and disease (Long et al., 2023; Meyer & North, 2019) – experts recommend accommodating expected shifts in species composition. Red fir forests are expected to experience substantial decline in the face of climate change (White et al., 2022).

- In areas that are transitioning out of being climatically suitable for red fir, consider favoring drought-tolerant species (60%), including rust-resistant sugar pine and Jeffrey pine, while still including a component of red fir (40%) in reforested stands.
- Specific site characteristics, at the project scale or units within the project, should also influence your selected species composition in the following ways:
 - **Topography:** When choosing to plant species other than red fir, plant a higher proportion of pine on ridgetops and south-facing slopes (North et al., 2009).
- **Following any drought or disease-driven die off**, you may also consider different species (e.g., Jeffrey pine) that are better adapted to higher elevation when near transition zone.



Climate Change Considerations

There are two important considerations with respect to planting for developing resilient forests in the face of climate change:

- **Wildfire resilience:** Wildfire resilience in mature stands can be promoted by planting trees at lower densities, which ultimately lowers fuel loads, and in variable arrangements to allow for the reintroduction of fire earlier. Continued management of fuels is still necessary to maintain wildfire resilience (see section below).
- **Drought resilience:** Competition for water resources between trees as they mature is lower at lower densities. Competition with other vegetation, especially shrubs, is lowered with clumped planting. Often, clumped stems will more quickly have interlocking crowns, which can shade out competing vegetation in the understory, as demonstrated in traditional, high-density plantings. Seedling clumps may also buffer conifer seedlings within the clump from the effects of future, lower severity fires.

Key Decisions

In addition to climate-related considerations, decisions about planting density and arrangement may reflect realities around survival of specific species and on certain sites. Survival within the initial establishment period can reflect differences in capacity for ongoing management and maintenance. Reforestation should generally not be undertaken if survival is anticipated to be low. In general, planting should be developed to plan for 80 – 90% survival of planted seedlings during establishment with some expected differences in planting survival across the landscape (e.g., lower survival is expected on xeric sites). Decisions around planting density and arrangement should reflect the abundance of natural regeneration present at the site. When available, use seedling plugs rather than bare root stock on harsher sites, for more sensitive seedlings, and during fall planting and periods of drought. Additionally, decisions around planting density and arrangement are dependent on capacity for post-planting management as described below.

<u>Density</u>

Variation in initial planting density recommendations reflect planting goals (e.g., desire to maximize merchantable timber, reestablish mature forest), capacity for follow-up treatments, and site characteristics. Experts recommend aiming for ~50-120 canopy trees per acre (TPA) with some age-class diversity in mature stands, depending on the forest type, site conditions, and long-term forest management goals (North et al., 2022; Safford & Stevens, 2017). While initial planting densities should assume that a percentage of seedlings planted will be lost to mortality, planting plans should include all available management actions that may improve establishment outcomes and support conservative seed use.

Which statement best reflects your post-planting management capacity and goals?



The capacity for post-planting thinning is high. There is a desire to maintain flexibility in planted tree management.



Higher Intensity Management

Some experts recommend initial planting densities of ~170-200 TPA. From these planting densities, final mature density targets can be reached with one or more rounds of pre-commercial thinning and natural causes of mortality. Decisions on specific levels of thinning should reflect climate and ecological conditions at the time. Planting at higher initial densities will require more extensive follow-up treatments, particularly to manage fuel loads to reduce wildfire risk and stand density to account for drought and insect driven mortality. See sections below about arrangement and postplanting management for more implications of higher density planting.



The capacity for post-planting thinning is low. There is a desire to minimize planted tree management.





Lower Intensity Management

Experts recommend an initial planting density closer to target levels for mature stands, ~60-150 TPA, with a buffer for anticipated mortality, recognizing that eventual seed production will increase future stand density as well. These lower initial planting densities will require a greater emphasis on use of microsites and more follow-up treatment to control competing vegetation. See section below about arrangement and post-planting management for more implications of lower density planting.

- **Following drought or disease-driven die off**, plant at the lower ends of the ranges listed above as these regions may be experiencing other issues (e.g., limited water availability and/or other stresses that can lead to an increased incidence of disease) that could also impact planted trees.
- There is less variation in recommended planting densities between mesic and xeric sites because trees are not competing with one another at the early stages. However, depending on the assessed capacity of the site, mesic sites can be planted with slightly higher initial densities than xeric sites.
- For red fir stands, if the stand was healthy prior to the disturbance, consider planting red fir in higher densities, as historic red fir stands were denser.
- **Topography**: Plant lower densities on ridgetops and south-facing slopes and higher densities on canyon bottoms and north-facing slopes within the ranges identified above.

<u>Arrangement</u>

Decisions about planting arrangement are interdependent with decisions related to planting density and capacity for follow-up treatments.

- Lower planting densities: Lower densities may be planted in a loosely gridded pattern with ample flexibility for use of microsites (Fig. 3). Other experts recommend planting with variable arrangement (e.g., clumping trees and leaving gaps) for ecological and operational reasons (Fig. 4). For example, post-planting management of shrubs is more effective and efficient when shrubs are concentrated in openings between clumped planting of trees. However, an increase in the complexity of planting pattern could slow the pace of implementation. Variable arrangements can include individual trees, clumps of trees, and openings (ICO) patterns.
 - Clusters may include 3-5 trees within a radius of 10-12 ft, with 20-25 ft between clusters, and openings of 1,000 sq ft, or a mixture of clusters and a gridded pattern.
 - Clusters should be located where microsite conditions can help ensure higher seedling survival.
 - Following drought or disease, retain clusters, but reduce the number of trees per cluster.
 - Prioritize planting clusters in areas with <60% shrub cover or where shrub cover is already patchy and heterogeneous.
- **Higher initial planting densities**: Some experts recommend planting in a roughly gridded pattern, allowing flexibility for utilization of microsites (Fig. 5). Through follow-up thinning and prescribed burns, a more heterogeneous forest structure can be cultivated while reducing stand density. Such higher densities are needed to meet industrial forestry objectives, since trees grown in a gridded pattern, as opposed to open grown trees, are often considered more desirable in form and implementation of subsequent maintenance treatments to reduce density is feasible and reliable. In these plantations, follow-up fuels reduction treatments are critical to address the increased risk from higher canopy bulk densities.

Variability in planting arrangement can also reduce future fire risk, allow for earlier reintroduction of fire through prescribed burning, and increase the likelihood of tree survival in extreme conditions.

- At the planting location, maximize use of shaded microsites (e.g., including rocks, live or dead shrubs, and snags, logs) to protect young seedlings from extreme heat, sun exposure, and soil drying (Fig. 2). When planting within a gridded pattern (Figs. 3 and 5), provide flexibility to adjust individual seedling location to take advantage of existing shaded microsites.
- At the project level, lower planting densities around the perimeter of a plantation can create a fuel break. This wide, lower density perimeter can be a focus for more intensive management of surface fuels. These could also be areas where a higher proportion of hardwoods are retained.
- In a planted fuel break, lower densities (75 TPA) may be planted to legal requirements (e.g., 24 x 24 ft spacing). Consider planting in variable planting arrangements to further introduce heterogeneity in fuel profile.

Timing

Existing guidance on the timing of planting is applicable for climate-informed reforestation, including preferring to plant in spring after snowmelt rather than fall planting because the soil moisture is high in the spring. However, it is worth noting the potential additional benefits of fall planting in the face of climate change. When planted in the fall, seedlings are ready to receive water by the spring even as the amount and timing of precipitation becomes more variable. While climatic variability may make fall freeze events more difficult to predict, seedlings planted in the fall will naturally harden off, preventing the need to freeze after lifting (taking the plant out of the nursery). The spring planting window between mid-April to June is restricted as snow melts and opens up access to higher elevation sites. The timing of spring is anticipated to shift with climate change, possibly leading to changes in the planting window. Additionally, planting as soon as possible following the initial disturbance remains a key factor leading to the success of any reforestation effort.



Planting with Microsites for Seedling Protection




Figure 2. Existing features within a planting area, such as shrub and herb cover, rocks, stumps, or downed logs, can create microsites with more shade, soil moisture, and protection from extreme heat and sun exposure. Planting to take advantage of these microsites can increase seedling growth and survival.



Lower Initial Planting Density in a Loose Grid



Figure 3. Lower density plantings can be arranged in a loose grid that allows flexibility for the use of microsites (Figure 2). Seedlings can be planted in higher densities on north-facing than south-facing slopes. This type of planting density and arrangement can maintain open areas for ecological diversity while supporting the development of a structured forest.



Lower Initial Planting Density with Variable Spacing




Figure 4. Lower density plantings can be spaced variably to create a site with individual seedlings, clusters of seedlings, and larger openings without any seedlings. Seedlings should be planted in areas where microsites can support their survival (Figure 2). Planting in clusters may offer advantages, as clumps of seedlings more quickly outcompete other vegetation and may be able to accept the reintroduction of beneficial fire earlier.



Higher Initial Planting Density in a Loose Grid




Figure 5. Higher density plantings arranged in a loose grid can support the growth of trees in a desirable form for merchantable timber and efficient implementation of post-planting thinning and maintenance treatments. Planting at higher densities in a gridded pattern should still allow flexibility for the use of microsites (Figure 2).

Key Actions

Reforestation does not end at planting. Continual management of forests following planting is required to ensure sustained climate and wildfire resilience into the future. As mentioned throughout this guidance, the capacity for varying degrees of post-planting treatment is a key driver of species, composition, density, and arrangement. A few common recommendations exist for all reforestation projects.

- Timely introduction of beneficial fire post-planting is a priority for climate-informed reforestation to
 promote wildfire resilience by reducing fuels, stand density, and reinforcing heterogeneity, and to
 create structural and composition diversity that promotes overall healthy forest ecosystems. It is also a
 very important ecosystem process that cannot be replaced or replicated with other treatment types.
 - Fire can be reasonably introduced 13-19 years after planting, depending on site productivity. At this point, a sufficient fuel bed has likely accumulated to carry fire, though studies are still in progress to refine these recommendations for when and how to reintroduce fire in reestablishing forests.
 - There are trade-offs when considering at which time of year to introduce fire. Fuel moisture is higher during spring and winter, which may produce a less intense fire and provide greater operational feasibility, including burn window. However, some early studies have shown higher mortality in spring burning. Fall burning would likely have less impact on the planted trees as they are not budding, is more aligned with a natural fire regime, and can result in greater surface fuels consumption within planted stands.
- Consider using one or more post-planting management practices (e.g., mechanical, hand treatments, fire, chemical, livestock grazing) timed appropriately to increase seedling vigor, reduce fuels, and increase reforestation success.
- Follow-up treatments to control competing vegetation, particularly aggressive shrub growth, is needed until trees are twice the height of the shrubs ("free to grow") to promote tree growth and subsequently to continue to address developing fuels.
- Use continued stand management practices to maintain densities within 20% of mature stand density target. Consider interplanting if densities drop significantly and extensively below the target, or to help create age class diversity in the stand.
- Post-planting monitoring is critical for assessing planting success (i.e., survival and growth of planted species, especially when planting a mixture of species from different seed zones), the effectiveness of site preparation actions to control competing vegetation, the need for density management, and the susceptibility to high-severity reburn based on fuel loads, and other management objectives (e.g., seedling spatial arrangement for facilitating heterogeneity in young stands).

Other post-planting treatment recommendations differ based on initial planting densities and arrangements as described above.



Lower Initial Planting Densities

- When planting at lower initial densities (~60-150 TPA), seedling survival may require additional postplanting treatments focused on controlling competing vegetation. Intensive shrub management can be reduced once trees are twice the height of shrubs, at which point shrub cover and tree cover provide for horizontal heterogeneity in fuel loading. However, additional treatments to reduce shrub cover may be necessary prior to reintroducing fire. There are many tools and methods available for this focus on managing understory growth (e.g., mastication, chemicals, fire, grazing), which can be selected based on existing guidance and operational considerations.
- In a low density planted site, use of prescribed fire may be more conservative since young tree mortality has a greater impact on reforestation outcomes with fewer trees. However, if the site has lower shrub cover and seedlings are planted in variable spacing, surface fuel loads and connectivity are likely lower, and use of prescribed fire can be more flexible. Additionally, lower density plantings may favor fall burning to avoid compromising a more limited seed source. Finally, early use of fire in low density plantings should be used primarily as a surface fuels management tool.



Higher Intensity Management

- Following higher initial planting densities (~170-200 TPA), there will be a greater need for focus on managing tree density including crown growth after the first few years. This management of crown growth and trees will require a greater focus on machinery and thinning. Post-planting management will entail multiple entries: (1) re-entry in early years 0-10 to reduce shrub cover, (2) thinning in the first 10-15 years to maintain crown separation, (3) multiple entries after 40 years, if a commercial thin is not already planned at this stage, to create gaps for regeneration of shade intolerant seedlings and to create diversity in age classes, as well as reducing fuels and inter-tree competition. Thinning following higher initial planting densities can be used to create heterogeneity in forest structure, and offers opportunities for altering species composition (e.g., prioritizing drought tolerant species), density, and disease management.
- Higher initial densities may allow for flexibility in managing future stand densities (e.g., trees could be removed later if site/microsite conditions are less productive than anticipated). However, higher densities also require follow-up treatments to reduce competition among neighboring stems and managing fuel loads.

• In higher density stands, which may allow for a higher tolerance of fire-killed trees, there is leeway to manage the stand with fire (i.e., pyrosilviculture) so long as fuel conditions are maintained to facilitate that option.



Other Contributors

Ben Blom, Christy Brigham, Brandon Collins, Michelle Coppoletta, Mark Egbert, Sarah Hennessy, Ryan Hilburn, Maurice Hunyh, Susie Kocher, Jonathan Long, Marc Meyer, Nicole Molinari, Malcolm North, Eric O'Kelley, Mark Pustejovsky, Thierry Rivard, Kristen Shive, Ryan Tompkins, Dana Walsh, Andrew Weinhart

Bibliography

- 1. Long, J. W., Walsh, D., Coppoletta, M., Tompkins, R. E., Meyer, M. D., Isbell, C., Bohlman, G. N., & North, M. P. (2023). Interventions to restore wildfire-altered forests in California (No. PSW-GTR-278; p. PSW-GTR-278). U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station. https://doi.org/10.2737/PSW-GTR-278
- 2. Lydersen, J. M., Collins, B. M., Coppoletta, M., Jaffe, M. R., Northrop, H., & Stephens, S. L. (2019). Fuel dynamics and reburn severity following high-severity fire in a Sierra Nevada, USA, mixed-conifer forest. Fire Ecology, 15(1). https://doi.org/10.1186/s42408-019-0060-x
- 3. Lydersen, J. M., & North, M. (2012). Topographic Variation in Structure of Mixed-Conifer Forests Under an Active-Fire Regime. Ecosystems, 15(7), 1134–1146. https://doi.org/10.1007/s10021-012-9573-8
- 4. Marsh, C., Blankinship, J. C., & Hurteau, M. D. (2023). Effects of nurse shrubs and biochar on planted conifer seedling survival and growth in a high-severity burn patch in New Mexico, USA. Forest Ecology and Management.
- 5. Marshall, L. A. E., Fornwalt, P. J., Stevens-Rumann, C. S., Rodman, K. C., Rhoades, C. C., Zimlinghaus, K., Chapman, T. B., & Schloegel, C. A. (2023). North-facing aspects, shade objects, and microtopographic depressions promote the survival
- and growth of tree seedlings planted after wildfire. Fire Ecology, 19(1). https://doi.org/10.1186/s42408-023-00181-8 6.Meyer, M. D., Long, J. W., & Safford, H. D. (2021). Postfire restoration framework for national forests in California (No. PSW-GTR-270); p. PSW-GTR-270). U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station. https://doi.org/10.2737/PSW-GTR-270
- 7. Meyer, M. D., & North, M. P. (2019). Natural Range of Variation of Red Fir and Subalpine Forests in the Sierra Nevada
- 8. Neuenkamp, L., Prober, S. M., Price, J. N., Zobel, M., & Standish, R. J. (2019). Benefits of mycorrhizal inoculation to ecological restoration depend on plant functional type, restoration context and time. Fungal Ecology, 40, 140-149.
- https://doi.org/10.1016/j.funeco.2018.05.004 9.North, M. P., Stevens, J. T., Greene, D. F., Coppoletta, M., Knapp, E. E., Latimer, A. M., Restaino, C. M., Tompkins, R. E., Welch, K. R., York, R. A., Young, D. J. N., Axelson, J. N., Buckley, T. N., Estes, B. L., Hager, R. N., Long, J. W., Meyer, M. D., Ostoja, S. M., Safford, H. D., ... Wyrsch, P. (2019). Tamm Review: Reforestation for resilience in dry western U.S. forests. Forest Ecology and Management, 432, 209-224. https://doi.org/10.1016/j.foreco.2018.09.007
- 10. North, M. P., Stine, P., O'Hara, K., Zielinski, W., & Stephens, S. (2009). An ecosystem management strategy for Sierran mixed-conifer forests. U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station.
- https://doi.org/10.2737/psw-gtr-220
 11. North, M. P., Tompkins, R. E., Bernal, A. A., Collins, B. M., Stephens, S. L., & York, R. A. (2022). Operational resilience in western US frequent-fire forests. Forest Ecology and Management, 507, 120004. https://doi.org/10.1016/j.foreco.2021.120004
- 12. Ritchie, M., Berrill, J.-P., Hammett, E., & Long, J. W. (2024). Early Responses to Crown Modification of California Black Oak Sprouts Initiated by High-Severity Wildfire. Journal of Forestry, 122(1), 70-78. https://doi.org/10.1093/jofore/fvad038
- 13. Safford, H. D., & Stevens, J. T. (2017). Natural range of variation for yellow pine and mixed-conifer forests in the Sierra Nevada, southern Cascades, and Modoc and Inyo National Forests, California, USA (No. PSW-GTR-256; p. PSW-GTR-256). U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station. https://doi.org/10.2737/PSW-GTR-256
- 14. Scott, J. H., & Burgan, R. E. (2005). Standard fire behavior fuel models: A comprehensive set for use with Rothermel's surface fire spread model (No. RMRS-GTR-153; p. RMRS-GTR-153). U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. https://doi.org/10.2737/RMRS-GTR-153
- 15. Sorenson, Q. M., Young, D. J. N., & Latimer, A. M. (2025). Tree planting outcomes after severe wildfire depend on climate,
- competition, and priority. Forest Ecology and Management, 575, 122346. https://doi.org/10.1016/j.foreco.2024.122346 16. Stewart, J. A. E., van Mantgem, P. J., Young, D. J. N., Shive, K. L., Preisler, H. K., Das, A. J., Stephenson, N. L., Keeley, J. E., Safford, H. D., Wright, M. Č., Welch, K. R., & Thorne, J. H. (2021). Effects of postfire climate and seed availability on postfire conifer regeneration. Ecological Applications, 31(3), e02280. https://doi.org/10.1002/eap.2280
- 17. White, A. M., Holland, T. G., Abelson, E. S., Kretchun, A., Maxwell, C. J., & Scheller, R. M. (2022). Simulating wildlife habitat dynamics over the next century to help inform best management strategies for biodiversity in the Lake Tahoe Basin, California. Ecology and Society, 27(2), art31. https://doi.org/10.5751/ES-13301-270231